

**PROBLEM 1 – 10 points**

You have three polarizers. Polarizer A has its transmission axis at  $0^\circ$  relative to the vertical; polarizer B has its transmission axis at  $30^\circ$  to the vertical; and polarizer C has its transmission axis at  $90^\circ$  to the vertical.

[4 points] (a) You can arrange the three polarizers in any order you wish. If the incident light is **unpolarized**, in what order should you place the three polarizers so as to maximize the intensity of the light emerging from the last polarizer in the sequence?

- Polarizer A, then B, then C       Polarizer C, then B, then A
- Either of the above, they'd give the same final intensity
- Polarizer B, then A, then C       Polarizer B, then C, then A
- It doesn't matter what order they're in, none of the wave gets through

Briefly justify your answer:

[3 points] (b) If the incident light is **polarized at  $30^\circ$  to the vertical**, aligned with the transmission axis of polarizer B, in what order should you place the three polarizers so as to maximize the intensity of the light emerging from the last polarizer in the sequence?

- Polarizer A, then B, then C       Polarizer C, then B, then A
- Either of the above, they'd give the same final intensity
- Polarizer B, then A, then C       Polarizer B, then C, then A
- It doesn't matter what order they're in, none of the wave gets through

[3 points] (c) In part (b), let's say the polarizers are arranged so that the light passes first through C, then through B, then through A. If the intensity of the light incident on the first polarizer is  $64 \text{ W/m}^2$ , find the intensity of the light emerging from:

- (i) polarizer C.
- (ii) polarizer B.
- (iii) polarizer A.

**PROBLEM 2 – 10 points**

A particular plane polarized electromagnetic wave, with a frequency of 100 MHz, is traveling through a vacuum in a direction we can call the  $x$ -axis. At  $t = 0$ , the electric field due to this wave at  $x = 0$  has a magnitude of 300 V/m.

[2 points] (a) What is the wavelength of this wave?

[2 points] (b) If this wave entered your eye, would you see anything? Explain why or why not.

[2 points] (c) At  $t = 0$  and  $x = 0$ , what is the magnitude of the magnetic field due to this wave?

[2 points] (d) How much time passes, after  $t = 0$ , before the electric and magnetic fields at  $x = 0$  are exactly the same as they are at  $t = 0$ ? State the minimum non-zero time.

[2 points] (e) If 300 V/m represents the amplitude of the electric field in this electromagnetic wave, what is the wave's average intensity?