

## Solutions and Spectroscopy Worksheet

As you work through the steps in the lab procedure, record your experimental values and the results on this worksheet. Use the exact values you record for your data to make later calculations.

## Part A - Determination of the Concentration of a Copper(II) Ion Solution

Complete the following table. (Enter concentrations to three significant figures.)

Data Table A. Calibration Curve of  $\text{Cu}^{2+}$  Solutions and Unknown

Stock $\text{Cu}^{2+}$ solution concentration _____ M						
Solution #	Target Volume of $\text{Cu}^{2+}$ , mL	Actual Volume of $\text{Cu}^{2+}$ , mL	Target Volume of $\text{H}_2\text{O}$ , ml	Actual Volume of $\text{H}_2\text{O}$ , ml	$[\text{Cu}^{2+}]$ , M (calculated)	Absorbance at ~620 nm (measured to 3 sf)
1	1.20		4.80			
2	2.40		3.60			
3	3.60		2.40			
4	4.80		1.20			
Equation of Trendline (to three significant figures):  $y = \text{_____} x + \text{_____}$						$R^2 = \text{_____}$ (to three significant figures)

Upload your graph as a file with a maximum size of 1 MB. (You will upload this file in the WebAssign question.)

Would you predict the absorbance of Solution 2 to be greater than or less than that of Solution 1?

Why?

What is the concentration of  $\text{Cu}^{2+}$  in your unknown solution? Record this concentration below. (*Hint*: Use the absorbance of the unknown and the trendline to solve for the  $\text{Cu}^{2+}$  concentration.)

Unknown #

Absorbance at 620 nm (measured)

$[\text{Cu}^{2+}]$  (calculated)

### Part B - Preparation of a Copper(II) ion Solution from Solid $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$

You desire to make a copper(II) solution at the same concentration as the unknown you just determined in Part A. How many grams of  $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$  are required to make 25.00 mL of this solution? Record the result as the target mass in Data Table B. (Enter concentrations to three significant figures.)

**Data Table B. Preparation of a  $\text{Cu}^{2+}$  Solution from solid  $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$**

Target $[\text{Cu}^{2+}]$ from Part A, <i>M</i>	Target Mass $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$ , g	Actual Mass $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$ , g	Absorbance of $\text{Cu}^{2+}$ solution at 620 nm	$[\text{Cu}^{2+}]$ calculated from absorbance, <i>M</i>

Would you predict the absorbance of your solution made from solid to be greater than or less than that of the unknown solution?

Why?

### Part C - Preparation of a Copper(II) Ion Solution by Dilution of a Stock $\text{CuSO}_4$ Solution

You desire to make a copper(II) solution at the same concentration as the unknown you determined in Part A. How many mL of the copper(II) stock solution are required to make 25.00 mL of this solution? Record the result as the target volume in Data Table C. (Enter concentrations to three significant figures.)

**Data Table C. Preparation of a  $\text{Cu}^{2+}$  Solution from stock  $\text{Cu}^{2+}$  Solution**

Stock $\text{Cu}^{2+}$ solution concentration _____ $M$				
Target $[\text{Cu}^{2+}]$ from Part A, $M$	Target Volume $\text{Cu}^{2+}$ solution, mL	Actual Volume $\text{Cu}^{2+}$ solution, mL	Absorbance of $\text{Cu}^{2+}$ solution at 620 nm	$[\text{Cu}^{2+}]$ calculated from absorbance, $M$

Would you predict the absorbance of your solution made from a dilution to be greater than or less than that of the unknown solution?

Why?